

Supplementary information for Hybridization Chain Reaction and Microscopy to Visualize Feeding Site– Specific Gene Expression in *Arabidopsis* Roots Infected by Plant-Parasitic Cyst Nematodes

Rahul S. Lathe¹, Anna Pijnacker¹, Norbert C.A. de Ruijter², Jose L. Lozano-Torres¹, Hendrik C. Korswagen^{3,4} and Christine W. Bruggeman^{3,*}

¹Laboratory of Nematology, Wageningen University & Research, Wageningen, The Netherlands

²Laboratory of Cell and Developmental Biology, Wageningen Light Microscopy Center, Wageningen University & Research, Wageningen, The Netherlands

³Hubrecht Institute, Utrecht, The Netherlands

⁴Division of Developmental Biology, Institute of Biodynamics and Biocomplexity, Utrecht University, Utrecht, the Netherlands

*For correspondence: c.bruggeman@hubrecht.eu




Figure S1. Single-molecule fluorescence in situ hybridization (smFISH) vs. hybridization chain reaction (HCR) spots in *Arabidopsis* seedling roots. Two independent samples stained with DAPI and with probes directed against *AtPP2A43*, either (A) using directly labeled probes (smFISH), imaged using a 60× objective, or (B) using probes with an adaptor to which fluorescently labeled hairpins are added in a second incubation step (HCR), imaged using a 40× objective. The HCR spots, though imaged with a smaller magnification objective, are clearly brighter than smFISH spots. Scale bars: 20 µm.